

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Meat and haem iron intake in relation to glioma in the European Prospective Investigation into Cancer and Nutrition study

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1766556> since 2021-01-13T09:20:22Z

Published version:

DOI:10.1097/CEJ.0000000000000331

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Published in final edited form as:

Eur J Cancer Prev. 2018 July ; 27(4): 379–383. doi:10.1097/CEJ.0000000000000331.

Meat and haem iron intake in relation to glioma in the European Prospective Investigation into Cancer and Nutrition study

Heather A. Ward¹, Alicia Gayle¹, Paula Jakszyn², Melissa Merritt¹, Beatrice Melin³, Heinz Freisling⁴, Elisabete Weiderpass^{5,6,7,8}, Anne Tjønneland⁹, Anja Olsen⁹, Christina C. Dahm¹⁰, Kim Overvad^{10,11}, Verena Katzke¹², Tilman Kühn¹², Heiner Boeing¹³, Antonia Trichopoulou^{14,15}, Pagona Lagiou^{14,15,16}, Andreas Kyrozi^{14,17}, Domenico Palli¹⁸, Vittorio Krogh¹⁹, Rosario Tumino²⁰, Fulvio Ricceri^{21,22}, Amalia Mattiello²³, H.Bas. Bueno-de-Mesquita^{1,24,25}, Petra H. Peeters^{1,26}, J.Ramón Quirós²⁷, Antonio Agudo²⁸, Miguel Rodríguez-Barranco^{29,30}, Nerea Larrañaga^{30,31}, José María Huerta^{30,32}, Aurelio Barricarte^{30,33,34}, Emily Sonestedt³⁵, Isabel Drake³⁵, Maria Sandström³, Ruth C. Travis³⁶, Pietro Ferrari⁴, Elio Riboli¹, and Amanda J. Cross¹

¹Department of Epidemiology and Biostatistics, Imperial College London, UK ²Unit of Nutrition, Environment and Cancer, Catalan Institute of Oncology, Barcelona – Spain ³Department of Radiation Sciences, Umeå University, Sweden ⁴Section of Nutrition and Metabolism, International Agency for Research on Cancer (IARC-WHO), Lyon, France ⁵Department of Community Medicine, Faculty of Health Sciences, University of Tromsø, The Arctic University of Norway, Tromsø, Norway ⁶Department of Research, Cancer Registry of Norway, Institute of population-based cancer research, Oslo, Norway ⁷Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden ⁸Genetic Epidemiology Group, Folkhälsan Research Center, Helsinki, Finland ⁹Diet, Genes and Environment, Danish Cancer Society Research Center, Copenhagen, Denmark ¹⁰Section for Epidemiology, Department of Public Health, Aarhus University, Denmark ¹¹Department of Cardiology, Aalborg University Hospital, Aalborg, Denmark ¹²Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany ¹³Department of Epidemiology, German Institute of Human Nutrition (DIfE), Potsdam-Rehbrücke, Germany ¹⁴Hellenic Health Foundation, Athens, Greece ¹⁵WHO Collaborating Center for Nutrition and Health, Unit of Nutritional Epidemiology and Nutrition in Public Health, Dept. of Hygiene, Epidemiology and Medical Statistics, University of Athens Medical School, Greece ¹⁶Department of Epidemiology, Harvard School of Public Health, Boston, USA ¹⁷Department of Neurology, Eginition Hospital, University of Athens Medical School, Greece ¹⁸Molecular and Nutritional Epidemiology Unit, Cancer Research and Prevention Institute (ISPO), Florence, Italy ¹⁹Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy

Corresponding author: Heather Ward: heather.ward@imperial.ac.uk.

Conflicts of interest: The authors declare that they do not have any competing interests.

Contribution of authors: AC originated the idea. AC was responsible for the study design and data management. HW and AG were responsible for the statistical analysis. AC, AG, and HW drafted the first version of the manuscript, contributed to interpretation of results and revised the manuscript critically for important intellectual content. All listed EPIC co-authors contributed data and reviewed the final manuscript.

Data sharing: For information on how to submit an application for gaining access to EPIC data and/or biospecimens, please follow the instructions at <http://epic.iarc.fr/access/index.php>

²⁰Cancer Registry and Histopathology Unit, “Civic- M.P.Arezzo” Hospital, ASP Ragusa, Italy ²¹Unit of Epidemiology, Regional Health Service, Grugliasco (TO), Italy ²²Unit of Cancer Epidemiology, Department of Medical Sciences, University of Turin, Turin, Italy ²³Dipartimento di Medicina Clinica e Chirurgia, Federico II University, Naples, Italy ²⁴Department for Determinants of Chronic Diseases (DCD), National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands ²⁵Department of Social & Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia ²⁶Department of Epidemiology, Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, the Netherlands ²⁷Public Health Directorate, Asturias, Spain ²⁸Unit of Nutrition and Cancer, Cancer Epidemiology Research Program. Catalan Institute of Oncology-IDIBELL. L'Hospitalet de Llobregat, Barcelona, Spain ²⁹Escuela Andaluza de Salud Pública. Instituto de Investigación Biosanitaria ibs.GRANADA, Hospitales Universitarios de Granada/Universidad de Granada, Granada, Spain ³⁰CIBER de Epidemiología y Salud Pública (CIBERESP), Spain ³¹Public Health Division of Gipuzkoa, Regional Government of the Basque Country, Spain ³²Department of Epidemiology, Murcia Regional Health Council, IMIB-Arrixaca, Murcia, Spain ³³Navarra Public Health Institute, Pamplona, Spain ³⁴Navarra Institute for Health Research (IdiSNA) Pamplona, Spain ³⁵Department of Clinical Sciences, Lund University, Malmö, Sweden ³⁶Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford UK

Abstract

Background—Diets high in red or processed meat have been positively associated with some cancers and several possible underlying mechanisms have been proposed, including iron-related pathways. However, the role of meat intake on adult glioma risk has yielded conflicting findings due to small sample sizes and heterogeneous tumor classifications.

Objective—The study objective was to examine red meat, processed meat and iron intake in relation to glioma risk in the European Prospective Investigation into Cancer and Nutrition (EPIC) study.

Methods—In this prospective cohort study, 408,751 individuals from nine European countries completed demographic and dietary questionnaires at recruitment. Multivariable Cox proportional hazards models were used to examine intake of red meat, processed meat, total dietary iron and haem iron in relation to incident glioma.

Results—During an average follow-up of 14.1 years, 688 incident glioma cases were diagnosed. There was no evidence that any of the meat variables (red, processed meat, or subtypes of meat) or iron (total or haem) were associated with glioma; results were unchanged when the first two years of follow-up was excluded.

Conclusion—This study suggests that there is no association between meat or iron intake and adult glioma. This is the largest prospective analysis of meat and iron in relation to glioma and as such provides a substantial contribution to a limited and inconsistent literature.

Keywords

EPIC; glioma; brain cancer; N-nitroso compounds; processed meat; red meat; dietary iron; haem iron

Introduction

Diets high in red or processed meat have been positively associated with several cancers (Abid et al., 2014), through complex mechanisms that include haem iron and the formation of *N*-nitroso compounds (NOCs) (Cross et al., 2003). The International Agency for Research on Cancer recently classified red meat and processed meat as class 2A and 1 carcinogens, respectively (Bouvard et al., 2015). Exposure to NOCs occurs from endogenous formation, which is directly related to red meat intake, and from exogenous exposure from nitrate- and nitrite-preserved meats. Haem iron, which is primarily found in red meat, has been shown to increase endogenous NOC formation (Cross, Pollock, & Bingham 2003) and may increase the risk of some common cancers (Fonseca-Nunes et al., 2014). NOCs may be highly carcinogenic and able to pass through the blood-brain barrier (McKinney, 2004), thus providing a mechanism through which the dietary components under examination could have been associated with glioma risk. A recent meta-analysis reported that processed meat may be positively associated with glioma (Wei et al., 2015); however, the majority of evidence available is from case-control studies, and iron intake has not been examined. Our aim was to examine intake of red meat, processed meat, total iron, and haem iron in relation to adult incident glioma risk in the large European Prospective Investigation into Cancer and Nutrition (EPIC) study.

Methods

From 1992-2000, the EPIC study recruited 521 448 adults (aged 25-70 years) from Denmark, France, Germany, Greece, Italy, Norway, the Netherlands, Spain, Sweden, and the United Kingdom (Riboli et al., 2002). Questionnaires on demographics, diet and other lifestyle factors were completed by participants at baseline. All EPIC centers measured height, weight, waist, and hip circumference except France, Norway, and Oxford where they were self-reported. The study was approved by all relevant ethical review boards, and all participants provided consent for the retention of acquired data and follow-up for incidence of cancer and death.

Dietary intake over the past year was assessed at baseline through country-specific questionnaires (Riboli, Hunt, Slimaniet al. 2002). Red and processed meat definitions within EPIC are described elsewhere (Rohrmann et al., 2013). The proportion of total iron that is haem iron was estimated to be 65% in cooked beef, 39% in pork and chicken, and 26% in fish (Balder et al., 2006).

Each participant was followed from baseline to cancer diagnosis, emigration, loss to follow-up, death, or end of study follow-up (which varied by center from June 2008-December 2013), whichever came first. Incident glioma cases were identified as site code C71 using the International Statistical Classification of Diseases and Related Health Problems (ICD-10) and morphology codes 9380-9460 from the International Classification of Diseases for Oncology (ICD-O).

France was not included in the present analysis as there were insufficient data to distinguish tumor histology (n=74,523). Among the remaining participants, we excluded those with

prevalent cancer at baseline (n=11,479), incomplete questionnaire data (n=6,837) or cancer follow-up data (n = 23,566) or if their energy intake versus energy requirement values were in the top or bottom 1% of the distribution (n=8,219). The analytical sample size was 408,751.

We examined energy-adjusted meat and iron as continuous variables and within fifths derived from the distribution among the total cohort; tests for linear trend were obtained using a continuous variable with values equal to the median intake within each fifth. Hazard ratios (HR) and 95% confidence intervals (CI) for glioma were estimated by Cox proportional hazards regression models, with person-years as the underlying time variable, stratified by age (rounded to one year), sex, and country. Smoking status, education, body mass index (kg/m²), physical activity, and intake of alcohol, total fat, saturated fat, fruit, vegetables and vitamin C were not included as covariates as none of them substantially (>10%) altered the risk estimates of the primary exposures. As sensitivity analyses, the first two years of follow-up were excluded to reduce the likelihood of dietary reports being influenced by potential changes in appetite related to undiagnosed glioma (reverse causality). The proportional hazards assumption was validated through examination of Martingale residuals.

Results

Descriptive characteristics of the cohort are presented in Table 1. During an average of 14.1 years of follow-up (5770398 person years), there were 324 glioma cases among men and 364 glioma cases among women.

Intake of red meat, processed meat, total iron, or haem iron was not associated with glioma (Table 2). These null results persisted for sex-specific dietary fifths, subtypes of meat and processed meat, after excluding incident glioma cases occurring within two years of recruitment (data not shown) and when absolute amounts of meat and iron were analyzed. No interactions were detected for age (above/below median) or smoking status for red meat, processed meat, or total iron (p values > 0.10). For haem iron, interactions were null by age (p 0.29). A marginally significant interaction by smoking status (p 0.08) suggested the potential for differences in the haem-glioma association by smoking status; however subgroup results are not provided given the lack of statistical significance.

Discussion

In this large prospective European study, red meat, processed meat, and total iron were not associated with the risk of glioma. The majority of previous studies to evaluate the NOC hypothesis in relation to glioma are retrospective studies; limited evidence supported the presence of an association (Lee et al., 1997). The results of the present study are consistent with other prospective results for red meat (Dubrow et al., 2010;Michaud et al., 2009)and processed meat (Wei, Zou, Cao et al. 2015); to the best of our knowledge, no other prospective investigations of total iron and haem iron in relation to glioma have been conducted.

The present study is the largest and only the fourth prospective cohort study to investigate the association between meat intake and adult glioma. Furthermore, it is the first study to assess this association in relation to total dietary iron intake. The prospective design precluded recall bias, and selection bias was minimized by the very high rate of follow-up over a long period of time. Equivalent baseline characteristics among exposure groups allowed for adequate comparisons to be made. The large number of incident cases of glioma was clearly defined and recorded which allowed for stratified analyses. A wide detailed range of meat intake was assessed in country-specific FFQs, which allowed for inclusion of a wide variety of dietary habits. A number of potential dietary and non-dietary factors were assessed for confounding, including measured height and weight for the majority of participants.

However, this study had several limitations. Food and nutrient intake was measured via multiple FFQs, which are subject to measurement error and may explain the lack of associations in the current study. However, the validation of these FFQs has been discussed previously and dietary estimates have been shown to be representative of European diets (Riboli, Hunt, Slimaniet al. 2002). In addition, direct estimates of dietary intake of individual NOCs were not calculated. Unrecorded information on cooking methods provided a limitation to explore other compounds related to meat such as heterocyclic amines or polycyclic aromatic hydrocarbons. Furthermore, usual dietary intake in the past year was assessed only at recruitment with no repeated measures, and may therefore not have captured diet during the etiologically relevant exposure period. Haem iron was calculated using specific factors for each type of meat, not measured directly, thus results are limited in interpretation due to assigning values using reported levels in the literature. In the future, the use of biomarkers of dietary intake may address some of limitations described above.

In summary, this study provides no indication that red meat or processed meat is associated with risk of adult glioma. Moreover, no significant associations were observed between iron intake and risk of glioma. These results do not provide support for the NOC hypothesis in relation to meat intake and adult glioma; however, future studies are warranted in order to confirm these findings and to determine whether specific biological mechanisms related to meat intake may be important.

Acknowledgments

The coordination of EPIC is financially supported by the European Commission (DG-SANCO) and the International Agency for Research on Cancer. The national cohorts are supported by Danish Cancer Society (Denmark); Ligue Contre le Cancer, Institut Gustave Roussy, Mutuelle Générale de l'Éducation Nationale, Institut National de la Santé et de la Recherche Médicale (INSERM) (France); German Cancer Aid, German Cancer Research Center (DKFZ), Federal Ministry of Education and Research (BMBF), Deutsche Krebshilfe, Deutsches Krebsforschungszentrum and Federal Ministry of Education and Research (Germany); the Hellenic Health Foundation (Greece); Associazione Italiana per la Ricerca sul Cancro-AIRC-Italy and National Research Council (Italy); Dutch Ministry of Public Health, Welfare and Sports (VWS), Netherlands Cancer Registry (NKR), LK Research Funds, Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund (WCRF), Statistics Netherlands (The Netherlands); ERC-2009-AdG 232997 and Nordforsk, Nordic Centre of Excellence programme on Food, Nutrition and Health (Norway); Health Research Fund (FIS), PI13/00061 to Granada, PI13/01162 to EPIC-Murcia, Regional Governments of Andalucía, Asturias, Basque Country, Murcia ((no. 6236) and Navarra, ISCIII RETIC (RD06/0020) (Spain); Swedish Cancer Society, Swedish Research Council and County Councils of Skåne and Västerbotten (Sweden); Cancer Research UK (14136 to EPIC-Norfolk; C570/A16491 and C8221/A19170 to EPIC-Oxford), Medical Research Council (1000143 to EPIC-Norfolk, MR/M012190/1 to EPIC-Oxford) (United Kingdom).

References

- Abid Z, Cross AJ, Sinha R. Meat, dairy, and cancer. *American Journal of Clinical Nutrition*. 2014; 100(1):386S–393S. available from: ISI:000337862200012. [PubMed: 24847855]
- Balder HF, Vogel J, Jansen MC, Weijenberg MP, van den Brandt PA, Westenbrink S, et al. Heme and chlorophyll intake and risk of colorectal cancer in the Netherlands cohort study. *Cancer EpidemiolBiomarkers Prev*. 2006; 15(4):717–725.
- Bouvard V, Loomis D, Guyton KZ, Grosse Y, Ghissassi FE, Benbrahim-Tallaa L, et al. Carcinogenicity of consumption of red and processed meat. *Lancet Oncol*. 2015; 16(16):1599–1600. [PubMed: 26514947]
- Cross AJ, Pollock JRA, Bingham SA. Haem, not protein or inorganic iron, is responsible for endogenous intestinal N-nitrosation arising from red meat. *Cancer Research*. 2003; 63(10):2358–2360. available from: ISI:000182981600003. [PubMed: 12750250]
- Dubrow R, Darefsky AS, Park Y, Mayne ST, Moore SC, Kilfoy B, et al. Dietary components related to N-nitroso compound formation: a prospective study of adult glioma. *Cancer EpidemiolBiomarkers Prev*. 2010; 19(7):1709–1722.
- Fonseca-Nunes A, Jakszyn P, Agudo A. Iron and cancer risk--a systematic review and meta-analysis of the epidemiological evidence. *Cancer EpidemiolBiomarkers Prev*. 2014; 23(1):12–31.
- Lee M, Wrensch M, Miike R. Dietary and tobacco risk factors for adult onset glioma in the San Francisco Bay Area (California, USA). *Cancer Causes Control*. 1997; 8(1):13–24. [PubMed: 9051318]
- McKinney PA. Brain tumours: Incidence, survival, and aetiology. *Journal of Neurology Neurosurgery and Psychiatry*. 2004; 75:12–17. available from: ISI:000225732900003.
- Michaud DS, Holick CN, Batchelor TT, Giovannucci E, Hunter DJ. Prospective study of meat intake and dietary nitrates, nitrites, and nitrosamines and risk of adult glioma. *AmJClinNutr*. 2009; 90(3):570–577.
- Riboli E, Hunt KJ, Slimani N, Ferrari P, Norat T, Fahey M, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr*. 2002; 5(6B):1113–1124. [PubMed: 12639222]
- Rohrmann S, Overvad K, Bueno-de-Mesquita HB, Jakobsen MU, Egeberg R, Tjønneland A, et al. Meat consumption and mortality--results from the European Prospective Investigation into Cancer and Nutrition. *BMC Med*. 2013; 11:63. [PubMed: 23497300]
- Wei YD, Zou DZ, Cao D, Xie P. Association between processed meat and red meat consumption and risk for glioma: A meta-analysis from 14 articles. *Nutrition*. 2015; 31(1):45–50. available from: ISI:000346685600006. [PubMed: 25466652]

Table 1

Baseline characteristics according to study-wide fifths of red meat, processed meat, total iron and haem iron intake

Characteristic ^a	Red meat intake			Processed meat intake			Total iron intake			Heme iron intake		
	Q1	Q3	Q5	Q1	Q3	Q5	Q1	Q3	Q5	Q1	Q3	Q5
No. of individuals	81748	81749	81751	81751	81751	81749	81748	81750	81751	81750	81751	81750
Age (years)	47.1	51.7	53.1	49.1	51.9	50.6	50.2	51.2	51.4	47.8	51.7	52.1
Height (cm)	166.9	166.0	167.8	164.8	167.2	168.1	168.7	166.9	164.5	166.9	166.3	167.4
BMI (kg/m ²)	24.5	26.0	26.7	25.5	25.8	26.3	25.2	25.6	26.8	24.2	26.0	26.9
Female (%)	71.9	67.8	55.1	71.3	64.2	59.0	59.2	67.0	67.0	73.4	67.7	52.7
Smoking status (%) ^b												
Never	54.9	45.3	38.6	55.0	44.4	40.5	42.2	46.0	50.3	53.6	44.6	41.9
Former	27.6	28.1	28.6	24.6	28.6	29.3	26.4	29.2	26.4	29.2	27.8	26.4
Current	16.3	24.9	31.9	18.5	25.7	28.8	29.8	23.6	21.8	15.7	26.0	30.7
Highest education level (%) ^b												
None	4.8	5.6	4.0	9.1	4.0	4.2	1.4	3.6	12.2	2.1	5.5	7.2
Primary school	16.3	31.0	33.6	22.8	29.2	31.0	30.5	26.9	29.5	15.0	30.5	35.2
Technical/professional school	23.3	25.8	29.7	17.9	28.1	30.5	28.0	27.6	20.9	24.2	27.5	25.1
Secondary school	16.7	16.1	13.8	15.7	16.4	13.8	18.6	16.2	12.1	17.0	16.4	13.5
Longer education	33.6	19.4	17.0	29.9	19.8	19.6	19.2	23.3	22.1	35.0	18.4	18.1
Physical activity (%) ^b												
Inactive	19.7	22.8	20.3	27.8	19.5	18.8	19.6	19.3	28.1	19.4	21.3	23.2
Moderately inactive	33.1	31.6	31.4	30.7	32.1	32.3	31.2	32.4	31.4	32.5	31.1	32.4
Moderately active	26.3	26.3	22.1	22.7	24.9	28.8	26.9	25.5	22.6	25.6	26.5	22.9
Active	19.5	17.2	22.9	17.6	21.1	17.7	19.5	20.7	16.8	20.5	18.6	19.9
Alcohol intake (g/day) ^c	3.7	4.9	8.6	3.7	6.0	5.1	3.6	6.3	4.9	4.1	5.2	6.9
Total fat intake (g/day) ^c	74.6	76.6	78.5	74.1	76.0	78.1	84.1	76.4	66.7	69.5	77.6	78.1
Saturated fat intake (g/day) ^c	28.5	28.4	29.4	24.9	29.1	30.3	34.5	29.2	21.9	26.4	29.0	28.4
Vitamin C intake (g/day) ^c	107.4	111.0	101.6	148.0	105.9	92.8	91.2	109.3	137.3	123.8	108.7	101.9
Vegetable intake (g/day) ^c	139.8	158.1	160.5	280.2	152.5	122.1	113.7	161.3	249.3	196.2	153.6	155.9
Fruit intake (g/day) ^c	180.5	192.7	161.0	276.1	183.1	125.2	144.7	186.7	234.8	212.6	185.5	162.3

^aValues are mean unless otherwise noted

^bData does not sum to 100% due to missing data

^cMedian values

Table 2

Meat and iron intake and the risk of incident glioma in the EPIC study

		Fifth of intake					P for trend
	Continuous ^c	1	2	3	4	5	
Red meat							
Intake, g/1000kcal ^a	20.2	2.67	10.18	17.36	26.52	44.46	
Cases	688	135	127	125	146	155	
HR ^b (95% CI)	1.01 (0.96,1.07)	1.00	1.00 (0.79,1.29)	1.02 (0.79,1.32)	0.99 (0.76,1.30)	0.99 (0.75,1.31)	0.93
Processed meat							
Intake, g/1000kcal ^a	15.3	1.11	6.48	12.28	19.68	36.89	
Cases	688	109	145	138	152	144	
HR ^b (95% CI)	1.02 (0.99,1.05)	1.00	0.99 (0.75,1.31)	1.04 (0.79,1.38)	1.02 (0.77,1.36)	1.12 (0.83,1.51)	0.36
Total Iron							
Intake, mg/1000kcal ^a	6.33	4.72	5.60	6.19	6.86	8.23	
Cases	688	133	142	149	143	121	
HR ^b (95% CI)	1.00 (0.94,1.07)	1.00	1.06 (0.84,1.35)	1.09 (0.85,1.38)	1.04 (0.81,1.34)	0.94 (0.71,1.24)	0.63
Heme Iron							
Intake, mg/1000kcal ^a	0.54	0.14	0.36	0.50	0.67	1.04	
Cases	688	134	114	144	154	142	
HR ^b (95% CI)	0.95 (0.74,1.22)	1.00	0.88 (0.68,1.15)	1.03 (0.80,1.35)	1.15 (0.90,1.49)	0.96 (0.73,1.26)	0.78

^aMedian intake.^bHazard ratios (HRs) and 95% confidence intervals (CI) from Cox proportional hazards models were adjusted for caloric intake (energy-adjusted quintiles), age, sex and country.^cRed meat intake (per 10g/day), processed meat intake (per 5g/day), total iron (per g/day), haem iron (per g/day)